

December, 2010

A quantum isomonodromy equation and its application to $\mathcal{N} = 2$ $SU(N)$ gauge theories

Yasuhiko Yamada

*Department of Mathematics, Faculty of Science
Kobe University, Hyogo, 657-8501, Japan*

Abstract

We give an explicit differential equation which is expected to determine the instanton partition function in the presence of the full surface operator in $\mathcal{N} = 2$ $SU(N)$ gauge theory. The differential equation arises as a quantization of a certain Hamiltonian system of isomonodromy type discovered by Fuji, Suzuki and Tsuda.

PACS: 11.25.HF, 11.15.-q.

Keywords: Quantum isomonodromy, Instanton partition function.

1 Introduction

In [1], Alday and Tachikawa formulated a combinatorial formula¹ for the instanton partition function Z_{inst} in the presence of the full surface operator in $\mathcal{N} = 2$ $SU(N)$ gauge theory, based on a result [2] about the affine Laumon space. Furthermore, they observed an interesting relation between Z_{inst} for $SU(2)$ theory and KZ equation with affine $SL(2)$ symmetry as an extension of the AGT relation [6]. A similar relation to Virasoro CFT including the irregular singularities was then examined in [3][4]. For $SU(N)$ case, the relation to affine $SL(N)$ conformal blocks was studied in [5].

¹We will recapitulate it in section 2, eq.(7).

In this note, we will study the differential equation satisfied by Z_{inst} from a point of view slightly different from KZ equations or W_N -algebras. Our basic strategy is to use the isomonodromy equations. It is known that some aspects of AGT relation have a natural interpretation [4][7][8] through the isomonodromy systems (or Painlevé equations for $SL(2)$ cases). That is, the $2d$ CFT's (=quantum isomonodromy systems) can be viewed as non-autonomous and quantum deformation of the Hitchin systems [9][10] (=Seiberg-Witten theory) arising through the Ω -deformation [11].

At the classical level, the relation between the Seiberg-Witten theory and isomonodromy equation is directly recognized by looking at the curves. For instance, the $SU(2)$ $N_f = 4$ Seiberg-Witten curve in Gaiotto form [12]

$$x^2 = \left(\frac{\mu_1}{z} + \frac{\mu_2}{z-1} + \frac{\mu_3}{z-t}\right)^2 - \frac{\kappa z + u}{z(z-1)(z-t)}, \quad \kappa = \mu_0^2 - \left(\sum_{i=1}^3 \mu_i\right)^2, \quad (1)$$

coincide with the Hamiltonian of the sixth Painlevé equation

$$H_{\text{VI}} = q(q-1)(q-t)p^2 - 2\{\mu_1(q-1)(q-1) + \mu_2q(q-t) + \mu_3q(q-1)\}p - \kappa q, \quad (2)$$

by the following variable change

$$p = x + \frac{\mu_1}{z} + \frac{\mu_2}{z-1} + \frac{\mu_3}{z-t}, \quad q = z, \quad H_{\text{VI}} = u. \quad (3)$$

Similar relations for degenerate cases were considered in [13].

We want to generalize this kind of correspondence to $SU(N)$ $N_f = 2N$ cases at the quantum level. The first problem is to look for the suitable isomonodromy system with higher rank symmetries. Fortunately, a nice candidate appeared in recent work [15][17]. The system, which we call Fuji-Suzuki-Tsuda (FST) equation², can be described as an isomonodromy defor-

²This equation (called P_{VI} -chain in [15]) was first considered by Tsuda in 2008, as a similarity reduction of his 'UC-hierarchy' (certain generalization of KP hierarchy). Independently, in the context of the Drinfeld-Sokolov hierarchy, it was obtained by Fuji-Suzuki [14] in case of $N = 3$ and generalized in [17]. Though our description in section 3 will follow the notation of [17], the isomonodromic picture considered here is naturally understood from the UC-hierarchy. Tsuda's construction contains also the Garnier type extension with spectral type $(N-1, 1), \dots, (N-1, 1), (1^N), (1^N)$ (see [16]).

mation of the $N \times N$ Fuchsian connection on \mathbb{P}^1 with regular singularity at $z = 0, 1, t, \infty$

$$D = \partial_z - \mathcal{A}, \quad \mathcal{A} = \frac{A_0}{z} + \frac{A_1}{z-1} + \frac{A_t}{z-t}, \quad (4)$$

with the following spectral type (= eigenvalue multiplicity of the residue matrices): A_0 and $A_\infty = -A_0 - A_1 - A_t$ are of type (1^N) while A_1 and A_t are of type $(N-1, 1)$ ³. We will recall the explicit form of the isomonodromy equation in section 3. Here, we look at the classical spectral curve in order to see its relation to the gauge theory.⁴ The curve for eq.(4) is given by

$$\begin{aligned} \det(v - z\mathcal{A}) &\propto \det\left((z-1)(z-t)(v - A_0) - z(z-t)A_1 - z(z-1)A_t\right) \\ &= \{(z-1)(z-t)\}^{N-1} f(z, v) = 0, \end{aligned} \quad (5)$$

where $v = z\partial_z$ is considered as a commuting variable. The factorization in the second line is due to the fact that A_1, A_t are of rank one. Hence $f(z, v)$ is of bi-degree $(2, N)$ in variables (z, v) and has the form

$$f(z, v) = t \prod_{i=1}^N (v - m_i) + z \left\{ \sum_{i=0}^{N-1} u_i v^i - (1+t)v^N \right\} + z^2 \prod_{i=1}^N (v - \tilde{m}_i), \quad (6)$$

where $A_0 \sim \text{diag}(m_1, \dots, m_N)$, $A_\infty \sim -\text{diag}(\tilde{m}_1, \dots, \tilde{m}_N)$. This is the desired form as the Seiberg-Witten curve for $SU(N)$ with $N_f = 2N$.

In the next section, we formulate our main conjecture that a differential equation (12) determines the instanton partition function Z_{inst} . The isomonodromic origin of our equation (12) is discussed in section 3.

2 Main conjecture

The instanton partition function $Z_{\text{inst}} = Z_{\text{inst}}(y; a, m, \tilde{m})$ in the presence of the full surface operator in $SU(N)$ $N_f = 2N$ superconformal gauge theory is

³Then, without loss of generality, one can assume that the eigenvalues of A_1 are $(0, \dots, 0, c)$ and similar for A_t .

⁴ Another indication comes from the special solutions. In both theories, generalized hypergeometric series ${}_N F_{N-1}$ appears [18][19], [20][21][22].

a function of N -variables $y = (y_1, \dots, y_N)$ depending on $3N - 1$ parameters $m = (m_1, \dots, m_N)$, $\tilde{m} = (\tilde{m}_1, \dots, \tilde{m}_N)$ and $a = (a_1, \dots, a_N)$, $\sum_{i=1}^N a_i = 0$.

Let us recall the combinatorial formula for Z_{inst} following [1][5]

$$Z_{\text{inst}} = \sum_{\lambda} Z(\lambda) \prod_{i=1}^N y_i^{k_i(\lambda)}. \quad (7)$$

Here the sum is taken over all N -tuples $\lambda = (\lambda^1, \dots, \lambda^N)$ of partitions $\lambda^i = (\lambda_1^i \geq \lambda_2^i \geq \lambda_3^i \geq \dots \geq \lambda_{\ell_i}^i > 0)$. The indices of $\lambda^i = (\lambda_j^i)$ will be extended to \mathbb{Z} by $\lambda^i = \lambda^{i+N}$ and $\lambda_j^i = 0$ ($j \leq 0$ or $j > \ell_i$). We put $|\lambda| = \sum_{i=1}^N |\lambda^i| = \sum_{i=1}^N \sum_{j \geq 1} \lambda_j^i$. The exponents $k_i(\lambda)$ are given by

$$k_i(\lambda) = \sum_{j \geq 1} \lambda_j^{i-j+1}. \quad (8)$$

The coefficients $Z(\lambda)$ are defined as

$$Z(\lambda) = \frac{n_f(a, \lambda, m) n_{\tilde{f}}(a, \lambda, \tilde{m})}{n_v(a, \lambda)}, \quad (9)$$

where

$$\begin{aligned} n_f(a, \lambda, m) &= n_{\text{bif}}(m, a, (\phi)^N, \lambda, 0), \\ n_{\tilde{f}}(a, \lambda, \tilde{m}) &= n_{\text{bif}}(a, \tilde{m}, \lambda, (\phi)^N, 0), \\ n_v(a, \lambda) &= n_{\text{bif}}(a, a, \lambda, \lambda, 0), \\ n_{\text{bif}}(a, b, \lambda, \mu, x) &= \prod_{t=1}^{|\lambda|+|\mu|} (w_t - x), \end{aligned} \quad (10)$$

and finally, the weights $w_t = w_t(a, b, \lambda, \mu)^5$ are determined by the formula [2]

⁵ $a, b \in \mathbb{C}^N$, and λ, μ are N -tuples of partitions. The constraints $\sum_{i=1}^N a_i = 0$, $\sum_{i=1}^N b_i = 0$ will be considered after the substitutions (10).

$$\begin{aligned}
\chi(a, b, \lambda, \mu) &= \sum_{t=1}^{|\lambda|+|\mu|} e^{w_t} \\
&= \sum_{k=1}^N \sum_{l' \geq 1} e^{a_k - b_{k-l'} + \epsilon_2(\lfloor \frac{l'-k}{N} \rfloor - \lfloor \frac{-k}{N} \rfloor)} \sum_{s=1}^{\mu_{l'}^{k-l'}} e^{\epsilon_1 s} \\
&\quad - \sum_{k=1}^N \sum_{l \geq 1} \sum_{l' \geq 1} e^{a_{k-l+1} - b_{k-l'} + \epsilon_2(\lfloor \frac{l'-k}{N} \rfloor - \lfloor \frac{l-k-1}{N} \rfloor)} (e^{\epsilon_1 \mu_{l'}^{k-l'}} - 1) \sum_{s=1}^{\lambda_l^{k-l+1}} e^{\epsilon_1(s - \lambda_l^{k-l+1})} \\
&\quad + \sum_{k=1}^N \sum_{l \geq 1} \sum_{l' \geq 1} e^{a_{k-l+1} - b_{k-l'+1} + \epsilon_2(\lfloor \frac{l'-k-1}{N} \rfloor - \lfloor \frac{l-k-1}{N} \rfloor)} (e^{\epsilon_1 \mu_{l'}^{k-l'+1}} - 1) \sum_{s=1}^{\lambda_l^{k-l+1}} e^{\epsilon_1(s - \lambda_l^{k-l+1})} \\
&\quad + \sum_{k=1}^N \sum_{l \geq 1} e^{a_{k-l+1} - b_k + \epsilon_2(\lfloor \frac{-k}{N} \rfloor - \lfloor \frac{l-k-1}{N} \rfloor)} \sum_{s=1}^{\lambda_l^{k-l+1}} e^{\epsilon_1(s - \lambda_l^{k-l+1})},
\end{aligned} \tag{11}$$

where $\lfloor x \rfloor$ is the largest integer such that $\lfloor x \rfloor \leq x$. The formula (11) is applicable to periodic parameters $a_{i+N} = a_i$, $b_{i+N} = b_i$. However, another convention is possible, where all terms $\epsilon_2(\lfloor \dots \rfloor - \lfloor \dots \rfloor)$ in the exponentials in eq.(11) are taken away in exchange for assuming quasi-periodicity $a_{i+N} = a_i + \epsilon_2$, $b_{i+N} = b_i + \epsilon_2$. In what follows, we will adopt the second option.

As compared with the above complicated formulae for Z_{inst} , the differential equation we propose is rather simple and defined as

$$\begin{aligned}
\mathcal{DZ}(y) &= \left\{ (\Delta_N + \sum_{i=1}^N u_i \vartheta_i) + (\prod_{j=1}^N y_j) (\Delta_N + \sum_{i=1}^N v_i \vartheta_i + \sum_{i=1}^N r_i s_i) \right. \\
&\quad \left. + \sum_{i=1}^N (y_i + y_i y_{i+1} + \dots + \prod_{j=0}^{N-2} y_{i+j}) (\vartheta_{i-1} - \vartheta_i + r_i) (\vartheta_{i-1} - \vartheta_i + s_i) \right\} \mathcal{Z} = 0,
\end{aligned} \tag{12}$$

where $y_{i+N} = y_i$, $\vartheta_i = y_i \frac{\partial}{\partial y_i}$, $\Delta_N = \frac{1}{2} \sum_{i=1}^N (\vartheta_i - \vartheta_{i+1})^2$. The parameters

u_i, v_i, r_i, s_i ($1 \leq i \leq N$) will be set as

$$\begin{aligned} u_i &= \frac{a_{i+1} - a_i}{\epsilon_1}, & v_i &= \frac{a_{i+1} - a_i + m_{i+1} - m_{i+2} + \tilde{m}_i - \tilde{m}_{i+1}}{\epsilon_1}, \\ r_i &= \frac{a_i - m_{i+1} - \epsilon_1}{\epsilon_1}, & s_i &= \frac{a_i - \tilde{m}_i}{\epsilon_1}, \end{aligned} \quad (13)$$

where $x_{i+N} = x_i + \epsilon_2$ for $x = a, m, \tilde{m}$ (while u_i, v_i, r_i, s_i are periodic).

The main claim in this note is the following

Conjecture. *The instanton partition function Z_{inst} is characterized as the unique formal power series solution of the form $\mathcal{Z} = 1 + \mathcal{O}(y)$ for the differential equation (12) with parameters (13).*

We have checked this conjecture for $N \leq 5$ up to total degree 5 in y -variables (in some cases by specializing the parameters to numerical values).

Under a degeneration limit $y_i \rightarrow \varepsilon^2 y_i$, $m_i(\tilde{m}_i) \rightarrow \varepsilon^{-1} \Lambda$, ($\varepsilon \rightarrow 0$), the differential equation (12) reduces to the Toda equation

$$\left(\Delta_N + \sum_{i=1}^N u_i \vartheta_i + \frac{\Lambda^2}{\epsilon_1^2} \sum_{i=1}^N y_i \right) \mathcal{Z} = 0, \quad (14)$$

whose relation to $\mathcal{N} = 2$ $SU(N)$ pure gauge theory has already been established by Braverman-Etingof [24] (see also [25][26][27]).

3 Origin of the differential equation

In this section, we explain an isomonodromic origin of our differential equation (12). As already mentioned in the introduction, the equation (12) is a quantization of the Fuji-Suzuki-Tsuda (FST) equation [14][15][17]. The FST equation can be written as a Hamiltonian system for $2(N-1)$ variables $q = (q_1, \dots, q_{N-1})$ and $p = (p_1, \dots, p_{N-1})$

$$t(t-1) \frac{dq_i}{dt} = \frac{\partial H}{\partial p_i}, \quad t(t-1) \frac{dp_i}{dt} = -\frac{\partial H}{\partial q_i}, \quad (15)$$

with parameters η and $\alpha = (\alpha_0, \dots, \alpha_{2N-1})$, $\sum_{j=0}^{2N-1} \alpha_j = 1$. The Hamiltonian $H = H(q, p, t; \eta, \alpha)$ is given by⁶

$$H = \sum_{i=1}^{N-1} H_{\text{VI}}(q_i, p_i; a_i, b_i, c_i, d_i) + \sum_{1 \leq i < j \leq N-1} (q_i - 1)(q_j - t) \{ (q_i p_i + \alpha_{2i-1}) p_j + p_i (q_j p_j + \alpha_{2j-1}) \}, \quad (16)$$

where H_{VI} is the Hamiltonian of sixth Painlevé equation

$$H_{\text{VI}}(q, p; a, b, c, d) = q(q-1)(q-t)p^2 - \{aq(q-1) + bq(q-t) + c(q-1)(q-t)\}p + dq, \quad (17)$$

$$\text{and } a_i = \sum_{j=i}^{N-1} \alpha_{2j}, \quad b_i = \sum_{j=0}^{i-1} \alpha_{2j}, \quad c_i = \sum_{j=0}^{N-1} \alpha_{2j+1} - \alpha_{2i-1} - \eta, \quad d_i = \eta \alpha_{2i-1}.$$

A quantization of the system is given by the Schrödinger equation⁷

$$\left\{ t(t-1) \frac{\partial}{\partial t} - H(q_i, \frac{\partial}{\partial q_i}) \right\} \tilde{\Psi}(q_1, \dots, q_{N-1}, t) = 0. \quad (18)$$

To connect this equation with Z_{inst} , we make a gauge transformation $\tilde{\Psi}(q, t) = y_1^{k_1} \dots y_N^{k_N} \Psi(y)$, together with the variables change

$$y_1 = q_1, \quad y_2 = \frac{q_2}{q_1}, \quad \dots, \quad y_{N-1} = \frac{q_{N-1}}{q_{N-2}}, \quad y_N = \frac{t}{q_{N-1}}, \quad (19)$$

$$q_i \frac{\partial}{\partial q_i} = \vartheta_i - \vartheta_{i+1} \quad (i = 1, \dots, N-1), \quad t \frac{\partial}{\partial t} = \vartheta_N,$$

($\vartheta_i = y_i \frac{\partial}{\partial y_i}$). Then the eq.(18) takes the form (in the following, we will

⁶This is a kind of coupled system of the Painlevé VI. Another equation of such type first found by Sasano [23] will also be important for some superconformal gauge theories.

⁷Here we will not consider the problem of operator ordering seriously since the ambiguities can be absorbed by shifts of parameters.

concentrate on $N = 3$ case)

$$\begin{aligned} \mathcal{D}'\Psi(y) = & \{\Delta_3 + u_1\vartheta_1 + u_2\vartheta_2 + u_3\vartheta_3 + M_0 \\ & + y_1(\vartheta_{31} + s_1)(\vartheta_{12} + r'_1) + y_2(\vartheta_{12} + s_2)(\vartheta_{23} + r'_2) + y_3(\vartheta_{23} + s_3)(\vartheta_{31} + r'_3) \\ & + y_2y_3(\vartheta_{12} + s_2)(\vartheta_{31} + r'_3) + y_3y_1(\vartheta_{23} + s_3)(\vartheta_{12} + r'_1) + y_1y_2(\vartheta_{31} + s_1)(\vartheta_{23} + r'_2) \\ & + y_1y_2y_3(\Delta_3 + v_1\vartheta_1 + v_2\vartheta_2 + v_3\vartheta_3 + M_1)\}\Psi(y) = 0, \end{aligned} \quad (20)$$

where $\vartheta_{ij} = \vartheta_i - \vartheta_j$, and $r'_i, s_i, u_i, v_i, M_0, M_1$ are some constants depending only on the parameters η, α and k_1, \dots, k_N (their precise expressions are not necessary). We can and will choose k_N so that $M_0 = 0$. Then, the equation (20) have a unique formal series solution of the form

$$\Psi(y) = \sum_{i,j,k=0}^{\infty} c_{ijk} y_1^i y_2^j y_3^k. \quad (c_{000} = 1) \quad (21)$$

The coefficients c_{ij0} are written in terms of the very-well-poised, balanced hypergeometric series

$$F(a_0; a_1, \dots, a_5) = \sum_{k=0}^{\infty} \frac{(a_0 + 2k)}{a_0} \prod_{i=0}^5 \frac{(a_i)_k}{(a_0 + 1 - a_i)_k}, \quad (22)$$

as

$$c_{ij0} = \frac{(r'_1 - j)_i (r'_2)_j (-s_1)_i (-s_2)_j}{i!j!(u_1 + 1 - j)_i (u_2 + 1)_j} F(u_1 - j; -i, -j, u_1 - r'_1 + 1, u_1 - s_2, -u_2 - j), \quad (23)$$

where $(x)_i = \Gamma(x + i)/\Gamma(x)$ is the Pochhammer symbol (see Appendix A for the proof). Similarly, c_{0ij} and c_{j0i} are given by cyclic shifts of parameters $x_i \rightarrow x_{i+1 \pmod{3}}$ ($x = r, s, u$).

On the other hand, corresponding coefficients of the instanton partition function

$$Z_{\text{inst}}(y) = \sum_{i,j,k=0}^{\infty} c_{ijk}^L y_1^i y_2^j y_3^k, \quad (24)$$

are obtained (at least for the first several terms) as

$$\begin{aligned}
c_{ij0}^L &= \sum_{k=0}^{\min(i,j)} Z(\{i, k\}, \{j - k\}, \phi) \\
&= (-1)^{i+j} \frac{\left(\frac{\epsilon_1 - a_1 + m_2}{\epsilon_1}\right)_i \left(\frac{\epsilon_1 - a_2 + m_3}{\epsilon_1}\right)_j \left(\frac{-a_1 + \tilde{m}_1}{\epsilon_1}\right)_i \left(\frac{-a_2 + \tilde{m}_2}{\epsilon_1}\right)_j}{i!j! \left(\frac{\epsilon_1 - \epsilon_1 j - a_1 + a_2}{\epsilon_1}\right)_i \left(\frac{\epsilon_1 - a_2 + a_3}{\epsilon_1}\right)_j} \\
&\quad \times F\left(\frac{-\epsilon_1 j - a_1 + a_2}{\epsilon_1}; -i, -j, \frac{-\epsilon_1 j + a_2 - a_3}{\epsilon_1}, \frac{\epsilon_1 - a_1 + m_3}{\epsilon_1}, \frac{-a_1 + \tilde{m}_2}{\epsilon_1}\right),
\end{aligned} \tag{25}$$

together with similar formulas for c_{0ij}^L , c_{j0i}^L obtained by shifts $x_1 \rightarrow x_2 \rightarrow x_3 \rightarrow x_1 + \epsilon_2$ ($x = a, m, \tilde{m}$). Comparing the coefficients c_{ij0} and c_{ij0}^L etc., we find that they almost coincide if we put

$$\begin{aligned}
r'_1 &= \frac{a_2 - m_3}{\epsilon_1}, \quad r'_2 = \frac{a_3 - (m_1 + \epsilon_2)}{\epsilon_1}, \quad r'_3 = \frac{a_1 - m_2}{\epsilon_1}, \\
s_1 &= \frac{a_1 - \tilde{m}_1}{\epsilon_1}, \quad s_2 = \frac{a_2 - \tilde{m}_2}{\epsilon_1}, \quad s_3 = \frac{a_3 - \tilde{m}_3}{\epsilon_1}, \\
u_1 &= \frac{a_2 - a_1}{\epsilon_1}, \quad u_2 = \frac{a_3 - a_2}{\epsilon_1}, \quad u_3 = \frac{(a_1 + \epsilon_2) - a_3}{\epsilon_1}.
\end{aligned} \tag{26}$$

In fact, under this parameter identification, the ratio of the coefficients c_{ij0}/c_{ij0}^L etc. are simply given by

$$\frac{c_{ijk}}{c_{ijk}^L} = (r'_1)_{i-j} (r'_2)_{j-k} (r'_3)_{k-i}. \tag{27}$$

Moreover, this relation (27) is satisfied also for $(ijk) = (111), (211), (121), (112)$ and further (as far as we checked), by putting

$$\begin{aligned}
M_1 &= (r'_3 - 1)s_1 + (r'_1 - 1)s_2 + (r'_2 - 1)s_3, \\
v_1 &= r'_1 - r'_3 - s_1 + s_2 - u_1, \quad v_2 = r'_2 - r'_1 - s_2 + s_3 - u_2, \\
v_3 &= r'_2 - r'_1 - s_3 + s_1 - u_3 - \frac{\epsilon_2}{\epsilon_1}.
\end{aligned} \tag{28}$$

In order to remove the factor in eq.(27), we make a "gauge transformation" (in momentum space) defined by

$$\mathcal{D}' \rightarrow \mathcal{D} = V^{-1} \mathcal{D}' V, \quad V = (r'_1)_{\vartheta_{12}} (r'_2)_{\vartheta_{23}} (r'_3)_{\vartheta_{31}}. \tag{29}$$

Under this transformation, the Euler derivatives remain invariant $\vartheta_i \rightarrow V^{-1}\vartheta_i V = \vartheta_i$. On the other hand, by using the relation $\vartheta_i y_j = y_j(\vartheta_i + \delta_{ij})$, the multiplication operators y_i are transformed as

$$\begin{aligned} y_1 &\rightarrow V^{-1}y_1V = y_1 \frac{(r'_1)\vartheta_{12}(r'_2)\vartheta_{23}(r'_3)\vartheta_{31}}{(r'_1)\vartheta_{12}+1(r'_2)\vartheta_{23}(r'_3)\vartheta_{31}-1} = y_1 \frac{\vartheta_{31}+r'_3-1}{\vartheta_{12}+r'_1}, \\ y_2 &\rightarrow y_2 \frac{\vartheta_{12}+r'_1-1}{\vartheta_{23}+r'_2}, \quad y_3 \rightarrow y_3 \frac{\vartheta_{23}+r'_2-1}{\vartheta_{31}+r'_3}, \end{aligned} \quad (30)$$

and hence

$$y_1 y_2 \rightarrow y_1 \frac{\vartheta_{31}+r'_3-1}{\vartheta_{12}+r'_1} y_2 \frac{\vartheta_{12}+r'_1-1}{\vartheta_{23}+r'_2} = y_1 y_2 \frac{\vartheta_{31}+r'_3-1}{\vartheta_{23}+r'_2}. \quad (31)$$

Then the differential operator \mathcal{D}' in eq.(20) is transformed into

$$\begin{aligned} \mathcal{D} &= V^{-1}\mathcal{D}'V = \Delta_3 + u_1\vartheta_1 + u_2\vartheta_2 + u_3\vartheta_3 \\ &+ y_1(\vartheta_{31}+s_1)(\vartheta_{31}+r_1) + y_2(\vartheta_{12}+s_2)(\vartheta_{12}+r_2) + y_3(\vartheta_{23}+s_3)(\vartheta_{23}+r_3) \\ &+ y_2 y_3(\vartheta_{12}+s_2)(\vartheta_{12}+r_2) + y_3 y_1(\vartheta_{23}+s_3)(\vartheta_{23}+r_3) + y_1 y_2(\vartheta_{31}+s_1)(\vartheta_{31}+r_1) \\ &+ y_1 y_2 y_3(\Delta_3 + v_1\vartheta_1 + v_2\vartheta_2 + v_3\vartheta_3 + M_1), \end{aligned} \quad (32)$$

where $(r_1, r_2, r_3) = (r'_3 - 1, r'_1 - 1, r'_2 - 1)$. Thus we arrived at the eq.(12) for $N = 3$ case.

4 Summary and discussions

In this note, we formulated an explicit differential equation (12) which is expected to determine the instanton partition function Z_{inst} in the presence of the full surface operator in $\mathcal{N} = 2$, $SU(N)$ gauge theory with $N_f = 2N$. The differential equation is derived as a quantization of the FST equation of isomonodromy type.

In [1][5], it was claimed that the partition function Z_{inst} is the conformal block of the affine Lie algebra SL_N (with the insertion of the K -operators). It is known [28][29] that the KZ equation satisfied by the conformal blocks can be interpreted as quantization of a typical isomonodromy system, the

Schlesinger equation. Hence, it is quite natural to expect a direct relation between the formulation of [1][5] and the isomonodromy approach here. For instance, the specialization of the primary fields V_χ , $\chi = \kappa\Lambda_1, \kappa\Lambda_{N-1}$ in [5] for the simple punctures agrees with the choice of the spectral type $(N-1, 1)$. More precise relations between these two formulations, in particular the understanding of the mysterious K -operators, will be an important future problem.

Though we have considered the isomonodromy deformation of an operator of the form (4), it can also be formulated by a scalar differential operator

$$L = \partial_z^N + u_2 \partial_z^{N-2} + \cdots + u_N. \quad (33)$$

Then the relation to W_N -algebras is also naturally expected (see [30][31] and references therein).

For the present, our understanding of the relation between $4d$ gauge theory and isomonodromy equation is still extrinsic. In [32] it was noted that the linear action of the loop operators (monodromy of surface operators) on the chiral partition function is independent of the gauge coupling. This observation may be a key ingredient for more conceptual understanding of the isomonodromic nature of gauge theories and the AGT relation.

Acknowledgments

We would like to thank H. Awata, H. Kanno, H. Nagoya, T. Suzuki, T. Tsuda and S. Yanagida for valuable discussions. The work is supported in part by JSPS grants-in-aid No.21340036 and No.S-19104002.

Appendix A : Proof of eq.(23)

From the equation (20), the coefficients $c_{i,j,0}$ in (21) are determined by

$$\begin{aligned} & c_{i-1,j,0}(i-1-j+r_1)(i-1-s_1) + c_{i-1,j-1,0}(j-1+r_2)(i-1-s_1) \\ & - c_{i,j-1,0}(j-1+r_2)(1+i-j+s_2) - c_{i,j,0}(i^2 - ij + j^2 + iu_1 + ju_2) = 0. \end{aligned} \quad (A.1)$$

Plugging (23) into this equation and rewriting the parameters as $a_0 = u_1 - j$, $a_1 = -i$, $a_2 = -j$, $a_3 = -u_2 - j$, $a_4 = u_1 - r'_1 + 1$, $a_5 = u_1 - s_2$, the relation we should prove reduces to

$$(a_0 a_1 - a_1^2 - a_2 a_3)F - (a_0 - a_1)a_1 F^{a_1} + \frac{(1+a_0)a_2 a_3 (1+a_0-a_1-a_4)(1+a_0-a_1-a_5)}{(1+a_0-a_1)(1+a_0-a_4)(1+a_0-a_5)} F^{a_0 a_2} + \frac{(1+a_0)a_1 a_2 a_3}{(1+a_0-a_4)(1+a_0-a_5)} F^{a_0 a_1 a_2 a_3} = 0, \quad (\text{A.2})$$

where $F = F(a_0; a_1, a_2, a_3, a_4, a_5)$ and $F^{a_0 a_2} = F|_{a_0 \rightarrow a_0+1, a_2 \rightarrow a_2+1}$ etc. Expanding the series F in each term, (A.2) can be written as

$$\sum_{k=0}^{\infty} \varphi_k \omega_k = 0, \quad (\text{A.3})$$

where $\varphi_k = \prod_{i=0}^k \frac{(a_i)_k}{(1+a_0-a_i)_k}$ and

$$\begin{aligned} \omega_k = & (a_0 a_1 - a_1^2 - a_2 a_3)(a_0 + 2k) - (a_0 - a_1 + k)(a_1 + k)(a_0 + 2k) \\ & + \frac{(1+a_0-a_1-a_4)(1+a_0-a_1-a_5)(a_0+k)(a_2+k)(a_3+k)(1+a_0+2k)}{(1+a_0-a_1+k)(1+a_0-a_4+k)(1+a_0-a_5+k)} \\ & + \frac{(a_0+k)(a_1+k)(a_2+k)(a_3+k)(1+a_0+2k)}{(1+a_0-a_4+k)(1+a_0-a_5+k)}. \end{aligned} \quad (\text{A.4})$$

Then the equation (A.3) follows from an identity

$$\varphi_k \omega_k = \varphi_{k+1} u_{k+1} - \varphi_k u_k, \quad u_k = k(k + a_0 - a_2)(k + a_0 - a_3), \quad (\text{A.5})$$

since the infinite sum is terminating: $\varphi_k = 0$ for $k > \min(i, j)$. \square

References

- [1] L. F. Alday and Y. Tachikawa, “Affine $SL(2)$ conformal blocks from $4d$ gauge theories,” arXiv:1005.4469.
- [2] B. Feigin, M. Finkelberg, A. Negut and R. Rybnikov, “Yangians and cohomology rings of Laumon spaces,” arXiv:0812.4656 [math.AG].
- [3] K. Maruyoshi and M. Taki, “Deformed prepotential, quantum integrable system and Liouville field theory,” arXiv:1006.4505.

- [4] H. Awata, H. Fuji, H. Kanno, M. Manabe and Y. Yamada, “Localization with a surface operator, irregular conformal blocks and open topological string,” arXiv:1008.0574.
- [5] C. Kozcaz, S. Pasquetti, F. Passerini and N. Wyllard, “Affine $sl(N)$ conformal blocks from $N = 2$ $SU(N)$ gauge theories,” arXiv:1008.1412.
- [6] L. F. Alday, D. Gaiotto and Y. Tachikawa, “Liouville correlation functions from four-dimensional gauge theories,” *Lett. Math. Phys.* **91** (2010) 167-197, arXiv:0906.3219.
- [7] J. Teschner, “Quantization of the Hitchin moduli spaces, Liouville theory, and the geometric Langlands correspondence I,” arXiv:1005.2846.
- [8] T.-S. Tai, “Uniformization, Calogero-Moser/Heun duality and Sutherland/bubbling pants,” arXiv:1008.4332.
- [9] A. Beilinson and V. Drinfeld, “Quantization of Hitchin’s integrable system and Hecke eigensheaves,” preprint (ca. 1995).
- [10] G. Bonelli and A. Tanzini, “Hitchin systems, $N = 2$ gauge theories and W -gravity,” arXiv:0909.4031.
- [11] N. Nekrasov and E. Witten, “The omega deformation, branes, integrability, and Liouville theory,” arXiv:1002.0888.
- [12] D. Gaiotto, “ $N = 2$ dualities,” arXiv:0904.2715.
- [13] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, “Cubic pencils and Painlevé Hamiltonians,” *Funkcial. Ekvac.* **48** (2005) 147-160, arXiv:nlin/0403009.
- [14] K. Fuji and T. Suzuki, “Drinfeld-Sokolov hierarchies of type A and fourth order Painlevé systems,” *Funkcial. Ekvac.* **53** (2010) 143-167, arXiv:0904.3434 [math-ph].
- [15] T. Tsuda, “From KP/UC hierarchies to Painlevé equations,” arXiv:1004.1347 [nlin.SI].
- [16] T. Tsuda, “UC hierarchy and monodromy preserving deformation,” Preprint: MI2010-7 (Kyushu University), arXiv:1007.3450 [math.CA].

- [17] T. Suzuki, “A class of higher order Painlevé systems arising from integrable hierarchies of type A ,” arXiv:1002.2685 [math.QA].
- [18] T. Suzuki, “A particular solution of a Painlevé system in terms of the hypergeometric function ${}_{n+1}F_n$,” SIGMA **6** (2010) 078, arXiv:1004.0059 [math-ph].
- [19] T. Tsuda, “Hypergeometric solution of a certain polynomial Hamiltonian system of isomonodromy type,” Quart. J. Math. (2010) 17pp, arXiv:1005.4130 [math.CA].
- [20] A. Mironov and A. Morozov, “On AGT relation in the case of $U(3)$,” Nucl. Phys. **B825** (2010) 1-37, arXiv:0908.2569.
- [21] R. Schiappa and N. Wyllard, “An A_r threesome: Matrix models, $2d$ CFTs and $4d$ $N = 2$ gauge theories,” arXiv:0911.5337.
- [22] H. Awata and Y. Yamada, “Five-dimensional AGT relation and the deformed β -ensemble,” Prog. Theor. Phys. **124** (2010) 227-262, arXiv:1004.5122.
- [23] Y. Sasano, “Higher order Painlevé equations of type $D_l^{(1)}$, RIMS Kokyuroku **1473** (2006) 143-163.
- [24] A. Braverman and P. Etingof, “Instanton counting via affine Lie algebras. II: From Whittaker vectors to the Seiberg-Witten prepotential,” arXiv:math/0409441.
- [25] B. Feigin, M. Finkelberg, I. Frenkel and R. Rybnikov, “Gelfand-Tsetlin algebras and cohomology rings of Laumon spaces,” arXiv:0806.0072 [math.AG].
- [26] A. Negut, “Laumon spaces and the Calogero-Sutherland integrable system,” Invent. Math. **178** (2009) 299-331, arXiv:0811.4454 [math.AG].
- [27] A. Braverman, B. Feigin, L. Rybnikov and M. Finkelberg, “A finite analog of the AGT relation I: finite W -algebras and quasimaps’ spaces,” arXiv:1008.3655 [math.AG].
- [28] N. Reshetikhin, “The Knizhnik-Zamolodchikov system as a deformation of the isomonodromy problem,” Lett. Math. Phys. **26** (1992), 167-177.

- [29] J. Harnad, “Quantum isomonodromic deformations and the Knizhnik-Zamolodchikov equations.” Symmetries and integrability of difference equations, 155-161, CRM Proc. Lecture Notes, 9, Amer. Math. Soc., Providence, RI, 1996.
- [30] S. Kanno, Y. Matsuo, S. Shiba and Y. Tachikawa, “ $N = 2$ gauge theories and degenerate fields of Toda theory,” arXiv:0911.4787.
- [31] N. Wyllard, “ W -algebras and surface operators in $N = 2$ gauge theories,” arXiv:1011.0289.
- [32] L. F. Alday, D. Gaiotto, S. Gukov, Y. Tachikawa and H. Verlinde, “Loop and surface operators in $N = 2$ gauge theory and Liouville modular geometry,” JHEP **1001**, 113 (2010), arXiv:0909.0945.